

On *Saaty's* and *Koczkodaj's* inconsistencies of pairwise comparison matrices¹

BOZÓKI, S. AND RAPCSÁK, T.

*Computer and Automation Research Institute,
Hungarian Academy of Sciences,
Budapest, Hungary*

Abstract

The aim of the paper is to obtain some theoretical and numerical properties of *Saaty's* and *Koczkodaj's* inconsistencies of pairwise comparison matrices (*PRM*). In the case of 3×3 *PRM*, a differentiable one-to-one correspondence is given between *Saaty's* inconsistency ratio and *Koczkodaj's* inconsistency index based on the elements of *PRM*. In order to make a comparison of *Saaty's* and *Koczkodaj's* inconsistencies for 4×4 pairwise comparison matrices, the average value of the maximal eigenvalues of randomly generated $n \times n$ *PRM* is formulated, the elements a_{ij} ($i < j$) of which were randomly chosen from the ratio scale

$$\frac{1}{M}, \frac{1}{M-1}, \dots, \frac{1}{2}, 1, 2, \dots, M-1, M,$$

with equal probability $1/(2M-1)$ and a_{ji} is defined as $1/a_{ij}$. By statistical analysis, the empirical distributions of the maximal eigenvalues of the *PRM* depending on the dimension number are obtained. As the dimension number increases, the shape of distributions gets similar to that of the normal ones. Finally, the inconsistency of asymmetry is dealt with, showing a different type of inconsistency.

1. Introduction

In multiattribute decision making (*MADM*), the aim is to rank a finite number of alternatives with respect to a finite number of attributes. Tender evaluations, public procurement processes, selections of applicants for positions, decisions on the best portfolios in investments are real-life decision situations in which *MADM* models can be used.

In solving a multiattribute decision problem, one needs to know the importances or weights of the not equally important attributes and also the evaluations of the alternatives with respect to the attributes. One technique, often used, is the method of pairwise comparisons a concept which is more than two hundred years old. *Condorcet* (1785) and *Borda* (1781) introduced it for voting problems in the 1780's by using only 0 and 1 in the pairwise comparison matrices. In experimental psychology, *Thorndike* (1920) and *Thurstone* (1927) used it in the 1920's. Especially, pairwise comparisons based on a ratio scale is one of the basic pillars of the *Analytic Hierarchy Process* (*Saaty*, 1980).

¹This research was supported, in part, by the Hungarian Scientific Research Fund, Grant Nos. OTKA-T043276, T043241 and K60480.

Given n objects, e.g., attributes or alternatives, we suppose that the decision maker(s) is (are) able to compare any two of them. In preference modelling, this assumption is called comparability. For any pairs (i, j) , $i, j = 1, 2, \dots, n$, the decision maker is requested to tell how many times the i -th object is preferred (or more important) than the j -th one, which result is denoted by a_{ij} .

By definition,

$$a_{ij} > 0; \quad (1.1)$$

$$a_{ii} = 1; \quad (1.2)$$

$$a_{ij} = \frac{1}{a_{ji}}, \quad (1.3)$$

for any pair of indices (i, j) , $i, j = 1, 2, \dots, n$. The name of matrices $\mathbf{A} = [a_{ij}]_{i,j=1,2,\dots,n} \in R^{n \times n}$ with properties (1.1-1.3) is pairwise comparison matrices or positive reciprocal matrices (*PRM*).

A pairwise comparison matrix \mathbf{A} is consistent if it satisfies the transitivity property

$$a_{ij}a_{jk} = a_{ik} \quad (1.4)$$

for any indices (i, j, k) , $i, j, k = 1, 2, \dots, n$. Otherwise, \mathbf{A} is inconsistent. It was shown by *Saaty* (1980) that a pairwise comparison matrix is consistent if and only if it is of rank one. When a pairwise comparison matrix \mathbf{A} is consistent, the normalized weights computed from \mathbf{A} are unique. Otherwise, an approximation of \mathbf{A} by a consistent matrix (determined by a vector) is needed.

A crucial point of this methodology is to determine the inconsistency of the pairwise comparison matrices. The only widely accepted rule of inconsistency is due to *Saaty* (1980), but his definition does not meet some important requirements (see *Section 2*). The aim of the paper is to make some comparison on *Saaty*'s and *Koczkodaj*'s inconsistencies of pairwise comparison matrices. The two approaches seem to be completely different, because while *Saaty*'s inconsistency ratio is an index for the departure from randomness, *Koczkodaj*'s inconsistency index is related to the departure from consistency with the possibility to locate inconsistency.

In *Section 2*, the question is how to investigate *Saaty*'s and *Koczkodaj*'s inconsistencies. In *Section 3*, the inconsistency formulas of 3×3 pairwise comparison matrices are studied from theoretical and computational points of view. A differentiable one-to-one correspondence is given between *Saaty*'s and *Koczkodaj*'s inconsistencies. In *Section 4*, by using statistical tools, the average value of the maximal eigenvalues of randomly generated $n \times n$ *PRM* is formulated, the elements a_{ij} ($i < j$) of which were randomly chosen from the ratio scale $\frac{1}{M}, \frac{1}{M-1}, \dots, \frac{1}{2}, 1, 2, \dots, M-1, M$, with equal probability $1/(2M-1)$ and a_{ji} is defined as $1/a_{ij}$. Then, a comparison of *Koczkodaj*'s inconsistency index and *Saaty*'s inconsistency ratio is given for 4×4 pairwise comparison matrices. In *Section 5*, the inconsistency of random pairwise comparison matrices is investigated and by statistical analysis, the empirical distributions of the maximal eigenvalues of the *PRM* depending on the dimension number are obtained. As the dimension number increases, the shape of distributions gets similar to that of the normal ones. In *Section 6*, the inconsistency of asymmetry is dealt with, showing a different type of inconsistency.

2. Inconsistency indices

In real-life decision problems, pairwise comparison matrices are rarely consistent. Nevertheless, decision makers are interested in the level of consistency of the judgements, which somehow expresses the goodness or “harmony” of pairwise comparisons totally, because inconsistent judgements may lead to senseless decisions.

Saaty (1980) proposed the following method for calculating inconsistency. Computing the largest eigenvalue λ_{max} of \mathbf{A} , he has shown that $\lambda_{max} \geq n$ and equals to n if and only if \mathbf{A} is consistent. Then, inconsistency index (CI_n) is defined by

$$CI_n = \frac{\lambda_{max} - n}{n - 1},$$

which gives the average inconsistency. Mathematically, inconsistency is not but a rescaling of the largest eigenvalue. Since $\lambda_{max} \geq n$, CI_n is always non-negative. The inconsistency index in its own has no meaning, unless we compare it with some benchmark to determine the magnitude of the deviation from consistency. Let a set of e.g., 500 random pairwise comparison matrices of size $n \times n$ be generated so that each element a_{ij} ($i < j$) be randomly chosen from the scale

$$\frac{1}{9}, \frac{1}{8}, \frac{1}{7}, \dots, \frac{1}{2}, 1, 2, \dots, 8, 9,$$

and a_{ji} is defined as $\frac{1}{a_{ij}}$. Let RI_n denote the average value of the randomly obtained inconsistency indices, which depends not only on n but on the method of generating random numbers, too. The inconsistency ratio (CR_n) of a given pairwise comparison matrix \mathbf{A} indicating inconsistency is defined by

$$CR_n = \frac{CI_n}{RI_n}.$$

If the matrix is consistent, then $\lambda_{max} = n$, so $CI_n = 0$ and $CR_n = 0$, as well. *Saaty* concluded that an inconsistency ratio of about 10% or less may be considered acceptable. The intuitive meaning of the 10 percent rule is skipped by several authors. A statistical interpretation of the 10 percent rule is given by *Vargas* (1982). More recently, *Saaty*’s threshold is 5% for 3×3 , and 8% for 4×4 matrices (*Saaty*, 1994).

It is emphasized that the inconsistency ratio CR_n is related to *Saaty*’s scale. The structuring process in AHP specifies that items to be compared should be within one order of magnitude. This helps avoid inaccuracy associated with cognitive overload as well as $a_{ij}a_{jk}$ relationships that are beyond the 1-9 scale, see e.g. *Lane* and *Verdini* (1989) and *Murphy* (1993). If only two attributes (or alternatives) are present, inconsistency is always zero, since the decision maker gives only one importance ratio.

Though the only one widely accepted rule of inconsistency for any order of matrix is due to *Saaty*, its consistency definition has some drawbacks. By *Koczkodaj* (1993), “*The author of this paper truly believes that failure of the pairwise comparison method to become more popular has its roots in the consistency definition.*” The major drawback of *Saaty*’s inconsistency definition seems to be the 10 percent rule of thumb. Another weakness of it is related to the location of inconsistency or rather its lack. Since an eigenvalue is a global characteristic of a matrix, by examining it, we cannot say which matrix element contributed to the increase of inconsistency. Some improvements can be found in *Saaty* (1990).

A general 3×3 pairwise comparison matrix has three comparisons a, b, c . In order to define *Koczkodaj's* inconsistency index (*Duszak and Koczkodaj (1994)* and *Koczkodaj (1993)*), consider a general 3×3 pairwise comparison matrix. Reduce this reciprocal matrix to a vector of three coordinates (a, b, c) . In the consistent cases, the equality $b = ac$ holds. It is always possible to produce three consistent reciprocal matrices (represented by three vectors) by computing one coordinate from the combination of the remaining two coordinates. These three vectors are: $\left(\frac{b}{c}, b, c\right)$, (a, ac, c) and $\left(a, b, \frac{b}{a}\right)$.

The inconsistency index of a general 3×3 pairwise comparison matrix is defined by *Koczkodaj* as the relative distance to the nearest consistent 3×3 pairwise comparison matrix represented by one of these three vectors.

Definition 2.1 *The inconsistency index of a general 3×3 pairwise comparison matrix is equal to*

$$CM(a, b, c) = \min \left\{ \frac{1}{a} \left| a - \frac{b}{c} \right|, \frac{1}{b} \left| b - ac \right|, \frac{1}{c} \left| c - \frac{b}{a} \right| \right\}. \quad (2.1)$$

The inconsistency index of an $n \times n$ ($n > 2$) reciprocal matrix \mathbf{A} is equal to

$$CM(\mathbf{A}) = \max \left\{ \min \left\{ \left| 1 - \frac{b}{ac} \right|, \left| 1 - \frac{ac}{b} \right| \right\} \quad \text{for each triad } (a, b, c) \text{ in } \mathbf{A} \right\}. \quad (2.2)$$

In the case of matrices of higher orders, the inconsistency index of a matrix element is equal to the maximum of CM of all possible triads which include this element.

Note that the inconsistency index is not a metric. By *Duszak and Koczkodaj (1994)*, the number of all possible triads of the $n \times n$ comparison matrices is equal to

$$n(n-1)(n-2)/3!. \quad (2.3)$$

In the case of 4×4 pairwise comparison matrices and a scale of 1 to 5, the threshold should be $1/3$ (*Koczkodaj et al., 1997*).

Other inconsistency indices have been introduced. The inverse inconsistency index suggested by *Dodd, Donegan and McMaster (1993)*, *Monsuur (1996)* applied a transformation of the maximal eigenvalues, *Peláez and Lamata (2003)* examined all the triples of elements and used the determinant to indicate consistency, furthermore, *Stein and Mizzi (2007)* obtained the harmonic consistency index. Another type of inconsistency index is the distance from a specific consistent matrix. *Chu, Kalaba and Spingarn (1979)* used the least squares estimation error, *Crawford and Williams (1985)* the logarithmic least squares estimation error, furthermore, *Aguarón and Moreno-Jiménez (2003)* the geometric consistency index for the logarithmic least squares method (the row geometric mean method).

Table 1 summarizes some weighting methods and inconsistency indices, namely, the eigenvector method (*EM*) and inconsistency ratio (*CR*) (*Saaty, 1980*), the least squares method (*LSM*) (*Chu, Kalaba and Spingarn, 1979*), the χ^2 squares method ($\chi^2 M$) (*Jensen, 1983*), the singular value decomposition method (*SVDM*) (*Gass and Rapcsák, 2004*) and *Koczkodaj's* inconsistency index (*Koczkodaj, 1993, 1994*), the logarithmic least squares method (*LLSM*), (*Crawford and Williams, 1985*) and *GCI*, (*Aguarón and Moreno-Jiménez, 2003*).

Method	The problem to be solved	Inconsistency (The optimal solution is denoted by \mathbf{w})	Threshold of acceptability
Eigenvector Method, <i>EM</i>	$\lambda_{max}\mathbf{w} = \mathbf{A}\mathbf{w},$ $\sum_{i=1}^n w_i = 1$	$CR_n = \frac{\lambda_{max}-n}{RI_n},$ where RI_n denotes the average CI value of $n \times n$ random matrices	$CR_n \leq 0.1$
Least Squares Method, <i>LSM</i>	$\min \sum_{i=1}^n \sum_{j=1}^n \left(a_{ij} - \frac{w_i}{w_j}\right)^2$ $\sum_{i=1}^n w_i = 1,$ $w_i > 0, \quad i = 1, 2, \dots, n$	$\sqrt{\sum_{i=1}^n \sum_{j=1}^n \left(a_{ij} - \frac{w_i^{LSM}}{w_j^{LSM}}\right)^2}$	
Chi Squares Method, $\chi^2 M$	$\min \sum_{i=1}^n \sum_{j=1}^n \frac{\left(a_{ij} - \frac{w_i}{w_j}\right)^2}{\frac{w_i}{w_j}}$ $\sum_{i=1}^n w_i = 1,$ $w_i > 0, \quad i = 1, 2, \dots, n$	$\sum_{i=1}^n \sum_{j=1}^n \frac{\left(a_{ij} - \frac{w_i^{\chi^2 M}}{w_j^{\chi^2 M}}\right)^2}{\frac{w_i^{\chi^2 M}}{w_j^{\chi^2 M}}}$	
Singular Value Decomposition Method, <i>SVDM</i>	$\mathbf{A}_{[1]} = \alpha_1 \mathbf{u}\mathbf{v}^T$ the best one rank approximation of \mathbf{A} in Frobenius norm; $w_i^{SVD} = \frac{u_i + \frac{1}{v_i}}{\sum_{j=1}^n u_j + \frac{1}{v_j}}$ $i = 1, 2, \dots, n$	$\sqrt{\sum_{i=1}^n \sum_{j=1}^n \left(a_{ij} - \frac{w_i^{SVD}}{w_j^{SVD}}\right)^2}$	$CM(\mathbf{A}) \leq 0.33$ $n = 4$ scale of $1, \dots, 5$
Logarithmic Least Squares Method, <i>LLSM</i>	$\min \sum_{i=1}^n \sum_{j=1}^n \left(\ln a_{ij} - \ln \frac{w_i}{w_j}\right)^2$ $\sum_{i=1}^n w_i = 1,$ $w_i > 0, \quad i = 1, 2, \dots, n$	$GCI(\mathbf{A}) = \frac{2 \sum_{i=1}^n \sum_{j=1}^n \left(\ln a_{ij} - \ln \frac{w_i^{LLSM}}{w_j^{LLSM}}\right)^2}{(n-1)(n-2)}$	

Table 1. Weighting methods and inconsistency indices

3. Inconsistency of 3×3 pairwise comparison matrices

In this part, it is shown that there exists a one-to-one correspondence between *Saaty's* inconsistency ratio and *Koczkodaj's* inconsistency index.

The general form of 3×3 positive reciprocal matrices is as follows:

$$\begin{pmatrix} 1 & a & b \\ 1/a & 1 & c \\ 1/b & 1/c & 1 \end{pmatrix}, \quad a, b, c \in R_+. \quad (3.1)$$

By *Tummala* and *Ling* (1998), the maximal eigenvalues of matrices (3.1) can be explicitly given by the function

$$\lambda_{\max}(a, b, c) = 1 + \sqrt[3]{\frac{b}{ac}} + \sqrt[3]{\frac{ac}{b}}, \quad (a, b, c) \in R_+^3. \quad (3.2)$$

A consequence of this formula is that λ_{\max} does not change if the elements a and b are multiplied by the same constant. Thus, the *CR*-inconsistencies of matrices

$$\begin{pmatrix} 1 & 2 & 2 \\ & 1 & 2 \\ & & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 7 & 7 \\ & 1 & 2 \\ & & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 9 & 9 \\ & 1 & 2 \\ & & 1 \end{pmatrix} \quad (3.3)$$

are equal, though the consistency violations in the matrices are different.

By formula (3.2), it is possible to make a connection between λ_{\max} and the inconsistency originated from the elements a, b, c of the positive reciprocal matrices.

Definition 3.1 In the case of (3.1), let T denote the maximum of two ratios, $\frac{ac}{b}$ and $\frac{b}{ac}$, i.e., $T = \max \left\{ \frac{ac}{b}, \frac{b}{ac} \right\}$.

If the matrix is consistent, T equals to 1, otherwise, $T > 1$.

Theorem 3.1 In the case of 3×3 pairwise comparison matrices, there exists a differentiable one-to-one correspondence for every pair of the inconsistency *CR* defined by *Saaty*, the inconsistency *CM* defined by *Koczkodaj* and $T = \max \left\{ \frac{ac}{b}, \frac{b}{ac} \right\}$ as follows:

$$CR(T) = \frac{\sqrt[3]{T} + \frac{1}{\sqrt[3]{T}} - 2}{2RI_3}, \quad T > 1. \quad (3.4)$$

$$T(CR) = \left(1 + RI_3 CR + \sqrt{RI_3 CR(2 + RI_3 CR)} \right)^3, \quad CR \in (0, \infty), \quad (3.5)$$

$$CM(T) = 1 - \frac{1}{T}, \quad T(CM) = \frac{1}{1 - CM}, \quad CM \in (0, 1), \quad (3.6)$$

$$CR(CM) = \frac{\frac{1}{\sqrt[3]{1 - CM}} + \sqrt[3]{1 - CM} - 2}{2RI_3}, \quad CM \in (0, 1), \quad (3.7)$$

$$CM(CR) = 1 - \frac{1}{\left(1 + RI_3 CR + \sqrt{RI_3 CR(2 + RI_3 CR)} \right)^3}, \quad CR \in (0, \infty). \quad (3.8)$$

Proof. From *Definition 3.1*, it follows that

$$\sqrt[3]{\frac{ac}{b}} + \sqrt[3]{\frac{b}{ac}} = \sqrt[3]{T} + \frac{1}{\sqrt[3]{T}}.$$

Since $\lambda_{\max} = 1 + \sqrt[3]{\frac{ac}{b}} + \sqrt[3]{\frac{b}{ac}}$, it can be written in the equivalent form

$$\lambda_{\max} = 1 + \sqrt[3]{T} + \frac{1}{\sqrt[3]{T}}. \quad (3.9)$$

Saaty defined the inconsistency ratio as $CR = \frac{\lambda_{\max} - n}{RI_n}$. Let us substitute $n = 3$ and (3.9) for the formula of CR , and (3.4) is proved.

Function $CR(T)$ is differentiable on the domain $T > 1$, and

$$CR'(T) = \frac{1 - \frac{1}{\sqrt[3]{T^2}}}{6RI_3\sqrt[3]{T^2}}, \quad (3.10)$$

which is positive if $T > 1$, consequently, CR is invertable in this domain. Its inverse function is equal to

$$T(CR) = \left(1 + RI_3 CR + \sqrt{RI_3 CR(2 + RI_3 CR)}\right)^3, \quad CR \in (0, \infty),$$

which proves (3.5).

Since

$$\begin{aligned} CM &= \min \left\{ \left| \frac{1}{a} \left| a - \frac{b}{c} \right| \right|, \left| \frac{1}{b} \left| b - ac \right| \right|, \left| \frac{1}{c} \left| c - \frac{b}{a} \right| \right| \right\} = \\ &= \min \left\{ \left| 1 - \frac{b}{ac} \right|, \left| 1 - \frac{ac}{b} \right|, \left| 1 - \frac{b}{ac} \right| \right\} = \min \left\{ \left| 1 - \frac{b}{ac} \right|, \left| 1 - \frac{ac}{b} \right| \right\}, \end{aligned}$$

it follows that

$$CM(T) = 1 - \frac{1}{T}, \quad CM'(T) = \frac{1}{T^2}, \quad T > 1,$$

and

$$T(CM) = \frac{1}{1 - CM}, \quad T'(CM) = \frac{1}{(1 - CM)^2}, \quad CM \in (0, 1).$$

In order to obtain $CR(CM)$, formulas (3.4) and (3.6) are used:

$$CR(CM) = \frac{\frac{1}{\sqrt[3]{1 - CM}} + \sqrt[3]{1 - CM} - 2}{2RI_3}, \quad CM \in (0, 1). \quad (3.11)$$

Similarly, formulas (3.5) and (3.6) are used to obtain

$$CM(CR) = 1 - \frac{1}{\left(1 + RI_3 CR + \sqrt{RI_3 CR(2 + RI_3 CR)}\right)^3}, \quad CR \in (0, \infty). \quad (3.12)$$

Since the derivatives

$$CR'(CM) = CR'(T) T'(CM) \quad \text{and}$$

$$CM'(CR) = CM'(T) T'(CR)$$

are different from zero, we have one-to-one correspondences. ■

Corollary 3.1 *In the case of 3×3 pairwise comparison matrices, the following properties are equivalent:*

$$CR \leq 10\%; \quad (3.13)$$

$$\frac{1}{2.63} = 0.38 \leq \frac{ac}{b} \leq 2.63; \quad (3.14)$$

$$CM \leq 0.62. \quad (3.15)$$

Proof. (3.13) \Leftrightarrow (3.14): Let $x = \sqrt[3]{\frac{b}{ac}}$. From (3.2) and since λ_{\max} corresponding to $CR = 10\%$ is 3.1048, (3.13) is equivalent to

$$x^2 - 2.1x + 1 \leq 0, \quad x > 0.$$

By solving equality $x^2 - 2.1x + 1 = 0, x > 0$, we obtain that $x_1^* \approx 1.38$ and $x_2^* = \frac{1}{x_1^*} \approx 0.7244$. Thus,

$$\frac{1}{x^*} \leq \sqrt[3]{\frac{b}{ac}} \leq x_1^*,$$

which is equivalent to the statement.

(3.13) \Leftrightarrow (3.15) follows from (3.11) and (3.12). ■

The intuitional meaning of (3.13) \Leftrightarrow (3.14) in *Theorem 3.1* may be interpreted by the following example. Let

$$\mathbf{A} = \begin{pmatrix} 1 & 2 & 6 \\ 1/2 & 1 & 3 \\ 1/6 & 1/3 & 1 \end{pmatrix}.$$

Now, $a = 2, b = 6, c = 3$, and \mathbf{A} is consistent ($\frac{ac}{b} = 1$). Let us fix a and b . If, e.g., $c = 4$, the inconsistency of matrix \mathbf{A} remains acceptable, because

$$\frac{ac}{b} = \frac{2 \cdot 4}{6} = 1.33 < 2.63.$$

The maximal value of c , for which matrix \mathbf{A} is acceptable by the 10% rule, is $3 \cdot 2.63 = 7.89$.

We remark that the CM -inconsistencies of matrices (3.3) are equal as well.

4. A comparison of *Saaty's* and *Koczkodaj's* inconsistency indices for 4×4 pairwise comparison matrices

Koczkodaj (1997) reported on concrete inconsistency index calculations based on a ratio scale $1/5, 1/4, 1/3, 1/2, 1, 2, 3, 4, 5$ for 4×4 pairwise comparison matrices. He remarked that in this case, an acceptable threshold of inconsistency is $1/3$. In order to make comparisons between *Saaty's* and *Koczkodaj's* inconsistency indices, we have to fit *Saaty's* threshold to the ratio scale $1/5, 1/4, 1/3, 1/2, 1, 2, 3, 4, 5$.

By the definition of CR , the rule of acceptability of a pairwise comparison matrix is that the maximal eigenvalue λ_{\max} should not be greater than a linear combination of the average λ_{\max} of randomly generated matrices, denoted by $\overline{\lambda_{\max}}$, with a coefficient 0.1, and $\lambda_{\max}(=n)$ of a consistent matrix, with a coefficient 0.9, i.e.,

$$CR \leq 0.10 \quad \Longleftrightarrow \quad \lambda_{\max} \leq 0.1\overline{\lambda_{\max}} + 0.9n. \quad (4.1)$$

We remark that $\overline{\lambda_{\max}}$ grows more rapidly (the slope of the approximating line is 2.76) than n .

Let $\bar{\lambda}_{\max}(n, M)$ denote the average value of the dominant eigenvalue of a randomly generated $n \times n$ matrix the elements of which are chosen from the ratio scale

$$\frac{1}{M}, \frac{1}{M-1}, \dots, \frac{1}{2}, 1, 2, \dots, M-1, M, \quad (4.2)$$

with equal probability $\frac{1}{2M-1}$.

Table 2 presents the values of $\bar{\lambda}_{\max}(n, M)$ for $n = 3, 4, \dots, 10$ and $M = 3, 4, \dots, 15$. $\bar{\lambda}_{\max}(n, M)$ can be well approximated by using a 4-parameter quasilinear regression.

Theorem 4.1

$$\bar{\lambda}_{\max} = 0.5625n - 0.621M + 0.2481Mn + 1.1478 + \varepsilon(n, M), \quad (4.3)$$

where $\varepsilon(n, M)$ denotes the approximation error of $\bar{\lambda}_{\max}(n, M)$.

Proof. The least-squares optimal solution of the 4-parameter quasilinear approximation problem

$$\bar{\lambda}_{\max}(n, M) \approx \alpha n + \beta M + \gamma nM + \delta$$

is as follows:

$$\begin{aligned} \alpha &= 0.5625, \\ \beta &= -0.6210, \\ \gamma &= 0.2481, \\ \delta &= 1.1478. \end{aligned}$$

The maximal approximate error $\varepsilon(n, M)$, while $3 \leq n \leq 10$, $3 \leq M \leq 15$, is 0.35. ■

The largest element (M)															
$\lambda_{\max}(n, M)$		3	4	5	6	7	8	9	10	11	12	13	14	15	
	3	3.236	3.369	3.505	3.641	3.778	3.913	4.049	4.184	4.317	4.450	4.582	4.714	4.845	
	4	4.555	4.884	5.226	5.578	5.933	6.292	6.652	7.015	7.378	7.742	8.106	8.472	8.834	
	5	5.901	6.445	7.017	7.606	8.208	8.819	9.435	10.057	10.683	11.311	11.940	12.574	13.209	
	6	7.256	8.019	8.823	9.656	10.506	11.370	12.245	13.128	14.017	14.913	15.811	16.712	17.620	
	7	8.615	9.597	10.633	11.705	12.801	13.915	15.045	16.185	17.331	18.486	19.645	20.810	21.978	
	8	9.977	11.177	12.442	13.752	15.091	16.452	17.830	19.220	20.620	22.028	23.445	24.865	26.290	
	9	11.339	12.757	14.252	15.797	17.377	18.983	20.605	22.244	23.895	25.555	27.222	28.896	30.574	
	10	12.702	14.337	16.059	17.840	19.658	21.504	23.373	25.258	27.155	29.063	30.980	32.903	34.832	

Table 2. Average value of the largest eigenvalues of random PRM depending on the largest element of the ratio scale

Let $CI(n, M)$, $RI(n, M)$ and $CR(n, M)$ denote the inconsistency index, the average value of the randomly obtained inconsistency indices and the inconsistency ratio with respect to the dimension number n and ratio scale (4.2), respectively. The theorem above provides an equivalent characterization of the 10 percent rule as follows:

Corollary 4.1

$$CR(n, M) = \frac{CI(n, M)}{RI(n, M)} \leq 0.10 \iff \lambda_{max} \leq 0.95625n - 0.0621M + 0.02481Mn + 0.1148. \quad (4.4)$$

Proof. By substituting (4.3) for (4.1), we have the result. ■

We emphasize that the condition for the acceptable inconsistency in (4.4) depends only on the data of the experimental pairwise comparison matrix, namely, on its dimension and its largest element. If we use a continuous ratio scale instead of the discrete scale by *Saaty*, the results remain almost the same.

The results of *Theorem 4.1* and *Corollary 4.1* can be used in the case of experimental pairwise comparison matrices. A set of 384 *PRM* taken from real-world AHP analyses were studied in *Gass and Standard* (2002). The experimental distribution of the numbers in the basic AHP comparison scale was unexpected. It seems that for these real-world problems, the decision makers did not use with large experimental probability the extreme comparison values of 8 and 9 (see *Table 1* in *Gass and Standard*, 2000). Consequently, in order to estimate the inconsistency more precisely, the influence of the pairwise comparisons determined by the decision makers can be taken into consideration through the largest ratio numbers, respectively.

Based on *Theorem 4.1* and *Table 3*, the inconsistency ratio $CR(4, 5)$ can be determined. By generating all the possible *PRM* ($9^6 = 531441$ matrices) with $CM \leq 1/3$ (1377 matrices) on the ratio scale $1/5, \dots, 1, \dots, 5$, *Figure 1* shows that the possible values of CM under $1/3$ are from the set $\{0, 1/6, 1/5, 1/4, 1/3\}$ and the total number of different pairs $(CM, CR(4, 5))$ is 14. We can state that the threshold $CM \leq 1/3$ corresponds to $CR(4, 5) \leq 0.0336$ (3.36%). It follows that *Koczkodaj's* inconsistency index for 4×4 pairwise comparison matrices with respect to ratio scale $1/5, \dots, 1, \dots, 5$, is stricter than that of *Saaty's*. It is noted that the 10% rule allows much higher CM -inconsistency when using the ratio scale $1/9, \dots, 1, \dots, 9$. An example is as follows:

$$\mathbf{A} = \begin{pmatrix} 1 & 1/8 & 2 & 6 \\ 8 & 1 & 7 & 9 \\ 1/2 & 1/7 & 1 & 2 \\ 1/6 & 1/9 & 1/2 & 1 \end{pmatrix},$$

where $CR = CR(4, 9) = 9.47\%$ and $CM = 0.8125$.




Figure 1. *Koczkodaj's* $CM \leq 1/3$ rule corresponds to $CR(4, 5) \leq 3.36\%$

It is emphasized that the threshold $CM \leq 1/3$ is given for 4×4 pairwise comparison matrices with respect to the ratio scale $1/5, \dots, 1, \dots, 5$. A question arises, namely, how to determine the threshold values for higher dimensions. A possible way is to use the “one grade off” or “two grades off” rules. By *Koczkodaj* (1997), “*An acceptable threshold of inconsistency is 0.33 because it means that one judgement is not more than two grades of the scale “different” from the remaining two judgements.*”

		The largest element (M)													
$RI(n, M)$		3	4	5	6	7	8	9	10	11	12	13	14	15	
	3	0.118	0.185	0.252	0.321	0.389	0.457	0.525	0.592	0.658	0.725	0.791	0.857	0.922	
	4	0.185	0.295	0.409	0.526	0.644	0.764	0.884	1.005	1.126	1.247	1.369	1.491	1.611	
	5	0.225	0.361	0.504	0.651	0.802	0.955	1.109	1.264	1.421	1.578	1.735	1.894	2.052	
Matrix	6	0.251	0.404	0.565	0.731	0.901	1.074	1.249	1.426	1.603	1.783	1.962	2.142	2.324	
size	7	0.269	0.433	0.606	0.784	0.967	1.153	1.341	1.531	1.722	1.914	2.108	2.302	2.496	
(n)	8	0.282	0.454	0.635	0.822	1.013	1.207	1.404	1.603	1.803	2.004	2.206	2.409	2.613	
	9	0.292	0.470	0.657	0.850	1.047	1.248	1.451	1.656	1.862	2.069	2.278	2.487	2.697	
	10	0.300	0.482	0.673	0.871	1.073	1.278	1.486	1.695	1.906	2.118	2.331	2.545	2.759	

Table 3. Average value of the inconsistency indices of random PRM depending on the largest element of the ratio scale

Let us consider the general form of 3×3 positive reciprocal matrices formulated in (3.1). In the consistent cases, $a = b/c$, $1/a = c/b$, $b = ac$, $1/b = 1/(ac)$, $c = b/a$, $1/c = a/b$. In the inconsistent cases, the approximation of an element by the other two elements can be considered by the grade difference

$$GD(a, b, c) = \min \left\{ \max \{ |a - b/c|, |1/a - c/b| \}, \right. \\ \left. \max \{ |b - ac|, |1/b - 1/(ac)| \}, \max \{ |c - b/a|, |1/c - a/b| \} \right\}.$$

Thus, the one grade off rule and the two grades off rule are

$$GD(a, b, c) \leq 1 \quad \text{and} \quad GD(a, b, c) \leq 2,$$

respectively.

In the case of matrices \mathbf{A} of higher orders, the one grade off rule and the two grades off rule (Koczkodaj *et al.*, 1997) are

$$GD(\mathbf{A}) = \max \left\{ GD(a, b, c) \quad \text{for each triad } (a, b, c) \text{ in } \mathbf{A} \right\} \leq 1 \quad \text{or} \quad 2.$$

Figure 2 shows that the threshold $CM \leq 1/3$ corresponds to $GD \leq 2/3$, which is close to the one grade off rule.




Figure 2. $CM \leq 1/3$ threshold corresponds to $GD \leq 2/3$

5. Inconsistency of random pairwise comparison matrices

Golden and *Wang* (1990) computed the random inconsistency indices and *Forman* (1990) the same for incomplete *PRM*. *Dodd*, *Donegan* and *McMaster* (1993) investigated the frequency distributions of random inconsistency indices and their statistical significance levels. *Lane* and *Verdini* (1989) determined the exact distribution of random inconsistency indices for 3×3 matrices, and random samples of 2500 matrices were produced and analysed for 4×4 to 10×10 and selected higher-order matrices, as well as stricter consistency requirements for 3×3 and 4×4 pairwise comparison matrices were suggested. *Standard* 2000 generated randomly 1000 *PRM*, but restricted the CR_n as follows. For $n = 3, 4$ or 5 , $CR_n < 0.1$ was required, for $n = 6$, $CR_n < 0.2$, and for $n = 7$, $CR_n < 0.3$. The computer was very slow in generating the random, low CR_n , *PRM* regarding sizes 6 and 7 and the results became more scattered as n increased. Additionally, regarding $n = 7$, there were no results for $CR_n < 0.1$. Due to these conditions, the low CR_n analysis was not run regarding matrices of sizes 8 and 9. A conclusion is that *Saaty*'s rule is statistically very strict for large *PRM*.

We have performed a statistical analysis of *CR* and *CM* inconsistencies. The aim of our simulation was to analyze the empirical distributions of the maximal eigenvalues λ_{max} of randomly generated pairwise comparison matrices. The elements a_{ij} ($i < j$) were randomly chosen from the scale

$$\frac{1}{9}, \frac{1}{8}, \frac{1}{7}, \dots, \frac{1}{2}, 1, 2, \dots, 8, 9,$$

and a_{ji} is defined as $\frac{1}{a_{ij}}$. In the paper, the assumption of equal probabilities is used.

In order to have equal probabilities ($\frac{1}{17}$), we used Matlab's *rand* function for simulating uniform distribution, the period of which is 2^{1492} . We have computed the average value of λ_{max} of randomly generated pairwise comparison matrices which is the basis of the mean random consistency index (RI_n). The values of λ_{max} corresponding to the $CR_n = 10\%$, the number of matrices which satisfies the $CR_n \leq 10\%$, $GD \leq 1$ and $GD \leq 2$ conditions were also computed. (It follows from the definition of CR_n that – if the comparisons are carried out randomly – the expected value of CR_n is 1.) In *Table 4*, n varies from 3 to 10, the sample size is 10^7 for all n .

In the case of 3×3 matrices, the sample size 10^7 is much larger than the number of different matrices $17^3 = 4913$. Thus, many (or all) of the matrices may have been counted more than once. The ratios of the numbers of matrices holding $CR \leq 10\%$, $GD \leq 1$ and $GD \leq 2$ compared to the sample size have been also computed if each matrix counted exactly once, and found to have almost the same results as above.

Our simulations are visualized in histograms, too. *Figures 3.a – 3.h* show the empirical distributions of λ_{max} on the lower horizontal axis and the corresponding consistency ratio CR_n on the upper horizontal axis. As n increases, the shape of distribution of λ_{max} gets similar to a normal one in our sample. For $n = 3$, a notable part of the randomly generated matrices satisfies the $CR_n \leq 10\%$ rule. The number of matrices with $CR_n \leq 10\%$ drastically decreases as n increases (see *Table 4*). Regarding $n = 8, 9, 10$, we have not found a matrix in the sample of ten million with acceptable inconsistency. Based on the results, it seems that the meaning of 10% for $n = 3$ is very different from $n = 8$, which is one of the weaknesses of the inconsistency ratio by *Saaty*. It is also interesting that *consistency* and *randomness* do not exclude each other: 1.7% of 3×3 random matrices (and 0.0014% of 4×4 random matrices) are consistent.

n	Sample size	Average value of λ_{\max}	RI_n	λ_{\max} corresponding to $CR = 10\%$	Number of matrices $CR \leq 10\%$	Number of matrices $GD \leq 1$	Number of matrices $GD \leq 2$
3	10^7	4.0484	0.5242	3.1048	2.08×10^6 1.42×10^6 with $GD \leq 1$ 2.0×10^6 with $GD \leq 2$	1.42×10^6 all with $CR \leq 10\%$	2.68×10^6 2.0×10^6 with $CR \leq 10\%$
4	10^7	6.6525	0.8842	4.265	3.15×10^5 2.76×10^4 with $GD \leq 1$ 1.55×10^5 with $GD \leq 2$	2.76×10^4 all with $CR \leq 10\%$	1.7×10^5 1.55×10^5 with $CR \leq 10\%$
5	10^7	9.4347	1.1087	5.4435	2.39×10^4 61 with $GD \leq 1$ 2371 with $GD \leq 2$	61 all with $CR \leq 10\%$	2404 2371 with $CR \leq 10\%$
6	10^7	12.244	1.2488	6.6244	770 0 with $GD \leq 1$ 13 with $GD \leq 2$	0 	13 all with $CR \leq 10\%$
7	10^7	15.045	1.3408	7.8045	9 0 with $GD \leq 1$ 0 with $GD \leq 2$	0 	0
8	10^7	17.831	1.4004	8.9831	0 	0 	0
9	10^7	20.604	1.4505	10.16	0 	0 	0
10	10^7	23.374	1.486	11.3374	0 	0 	0

Table 4. Average value of λ_{max} of randomly generated pairwise comparison matrices, RI_n , the number of matrices with $CR \leq 10\%$, $GD \leq 1$ and $GD \leq 2$

6. Inconsistency of asymmetry

A conceptual weakness of some weighting method is related to the issue of asymmetry. The question: “To what extent does alternative i dominate j ?” may be replaced by the question “To what extent is j dominated by i ?” The answers to these questions are logically reciprocal. If a technique is applied first to the pairwise comparison matrix \mathbf{A} , yielding a solution \mathbf{w} , and then to the transpose \mathbf{A}^T , yielding a solution \mathbf{w}' , is $\frac{w_i}{w_j} = \frac{w'_j}{w'_i}$ for every pair (i, j) ?

EM does not possess this asymmetry property, since the principal right and left eigenvectors of \mathbf{A} are not elementwise reciprocal in the cases of inconsistent pairwise comparison matrices. Consequently, a conceptual limitation of EM is the lack of asymmetry with respect to \mathbf{A} and \mathbf{A}^T , which means that, for $n \geq 4$, there exist, generally, two competing solutions (*Johnson et al.*, 1979). Now, it will be shown that the property of asymmetry is related to the inconsistency.

Definition 6.1 Let \mathbf{A} be a pairwise comparison matrix, \mathbf{w} and \mathbf{w}' the priority vectors of \mathbf{A} and \mathbf{A}^T , respectively. The invariance under transpose holds if

$$w_i \geq w_j \quad \text{implies} \quad w'_i \leq w'_j, \quad \forall(i, j), \quad i, j = 1, \dots, n. \quad (6.1)$$

It follows from the definitions that LSM , χ^2M and $LLSM$ defined in *Table 1* always fulfil the property of invariance under transpose. $SVDM$ takes this asymmetry, in some sense, into account.

Lemma 6.1 $SVDM$ fulfils the invariance under transpose if and only if

$$\frac{u_i v_i + 1}{u_j v_j + 1} \geq \frac{v_i}{v_j} \quad \text{implies} \quad \frac{u_i v_i + 1}{u_j v_j + 1} \leq \frac{u_i}{u_j}, \quad \forall(i, j) \quad i, j = 1, \dots, n, \quad (6.2)$$

where \mathbf{u} and \mathbf{v} are the left and right singular vectors belonging to the largest singular value of \mathbf{A} , respectively.

Proof. By the formula in *Table 1*, the invariance under transpose holds if and only if

$$u_i + \frac{1}{v_i} \geq u_j + \frac{1}{v_j} \quad \text{implies} \quad v_i + \frac{1}{u_i} \leq v_j + \frac{1}{u_j}, \quad \forall(i, j) \quad i, j = 1, \dots, n,$$

which is equivalent to

$$\frac{u_i v_i + 1}{u_j v_j + 1} \geq \frac{v_i}{v_j} \quad \text{implies} \quad \frac{u_i v_i + 1}{u_j v_j + 1} \leq \frac{u_i}{u_j}, \quad \forall(i, j) \quad i, j = 1, \dots, n. \quad \blacksquare$$

10^8 matrices of size 5×5 have been generated randomly in order to detect the rank reversals of the weights computed from the left and right eigenvectors. Based on our hypothesis, the frequency of rank reversals varies as the CR inconsistency ratio changes. By *Table 5* and *Figure 4*, the frequency of rank reversals increases as the CR increases. We can conclude that the larger the CR -inconsistency is, the more often the EM violates the property of invariance under transpose. Since no “cut off” point appears in *Figure 4*, this seems to be another reason for reconsidering the asymmetry property.

The next example (*Dodd et al.*, 1995) shows that a good inconsistency ratio CR does not exclude the rank reversal between the weights computed from the left and right eigenvectors. Let

$$\mathbf{A} = \begin{pmatrix} 1 & 1 & 3 & 9 & 9 \\ 1 & 1 & 5 & 8 & 5 \\ 1/3 & 1/5 & 1 & 9 & 5 \\ 1/9 & 1/8 & 1/9 & 1 & 1 \\ 1/9 & 1/5 & 1/5 & 1 & 1 \end{pmatrix},$$

where $CR(\mathbf{A}) = 0.0820$, the weights of the right eigenvector

$$\mathbf{w}^T = (36.5652, 38.9564, 16.7155, 3.4693, 4.2936),$$

and the weights of the left eigenvector

$$\mathbf{w}'^T = (40.6431, 36.4208, 15.0669, 3.4391, 4.4302).$$

It is interesting that $GD(\mathbf{A}) = 4.1111$. There remain open questions, namely, how to detect and eliminate the inconsistency of asymmetry.

Levels of inconsistency ratio CR	Number of rank reversals of the weight vectors corresponding to the left and right eigenvectors	Number of matrices	Frequency of rank reversals
$CR \leq 0.01$	8	162	0.049
$0.01 < CR \leq 0.02$	81	1138	0.071
$0.02 < CR \leq 0.03$	288	3414	0.084
$0.03 < CR \leq 0.04$	685	7130	0.096
$0.04 < CR \leq 0.05$	1253	12645	0.099
$0.05 < CR \leq 0.06$	2096	19827	0.106
$0.06 < CR \leq 0.07$	3342	29686	0.113
$0.07 < CR \leq 0.08$	5284	41400	0.128
$0.08 < CR \leq 0.09$	7896	55105	0.143
$0.09 < CR \leq 0.10$	10819	70885	0.153
$0.10 < CR \leq 0.11$	14371	88104	0.163
$0.11 < CR \leq 0.12$	18743	1.07×10^5	0.174
$0.12 < CR \leq 0.13$	23362	1.28×10^5	0.182
$0.13 < CR \leq 0.14$	27841	1.50×10^5	0.185
$0.14 < CR \leq 0.15$	33402	1.73×10^5	0.193
$0.15 < CR \leq 0.16$	39344	1.97×10^5	0.199
$0.16 < CR \leq 0.17$	44851	2.21×10^5	0.203
$0.17 < CR \leq 0.18$	50847	2.46×10^5	0.207
$0.18 < CR \leq 0.19$	57625	2.69×10^5	0.214
$0.19 < CR$	not analysed	9.82×10^7	not analysed

Table 5. Frequency of rank reversals of the weight vectors corresponding to the left and right eigenvectors with respect to different levels of inconsistency ratio CR




Figure 4. Frequency of rank reversals of the weight vectors corresponding to the left and right eigenvectors with respect to different levels of inconsistency ratio CR

7. Concluding remarks

In the paper, some theoretical and numerical properties of *Saaty's* and *Koczkodaj's* inconsistencies of *PRM* are investigated. Based on the results, it seems that the determination of the inconsistency of *PRM* has some drawbacks, thus the improvement of the notion of inconsistency should be necessary.

Related to *Saaty's* inconsistency ratio, some basic questions are as follows:

What is the relation between an empirical matrix from human judgements and a randomly generated one? Is an index obtained from several hundreds of randomly generated matrices the right reference point for determining the level of inconsistency of pairwise comparison matrix built up from human decisions, for a real decision problem? How to take the size of matrices into account in a more precise form?

Related to *Koczkodaj's* consistency index, a major question seems to be the elaboration of the thresholds in higher dimensions or to replace the index by a refined grade off rule.

The existence of the inconsistency of asymmetry shows the complexity of the problem. By the example in *Section 6*, *Saaty's* consistency of *PRM* is insufficient to exclude asymmetric inconsistency, therefore, this latter should be considered as a separate issue. Thus, it seems that only one inconsistency index is insufficient for describing the inconsistency.

Acknowledgement

The authors thank the Referees for the remarks and advice. The distinction of *Saaty's* and *Koczkodaj's* inconsistencies, given by one of the Referees, is emphasized.

References

- Aguarón, J. and Moreno-Jiménez, J.M., The geometric consistency index: Approximated thresholds, *European Journal of Operational Research* 147 (2003) 137-145.
- Borda, J.C. de, Mémoire sur les élections au scrutin, *Histoire de l'Académie Royale des Sciences*, Paris, 1781.
- Chu, A.T.W., Kalaba, R.E. and Spingarn, K., A comparison of two methods for determining the weight belonging to fuzzy sets, *Journal of Optimization Theory and Applications* 4 (1979) 531-538.
- Condorcet, M., Essai sur l'application de l'analyse à la probabilité des décisions rendues à la pluralité des voix, Paris, 1785.
- Crawford, G. and Williams, C., A note on the analysis of subjective judgment matrices, *Journal of Mathematical Psychology* 29 (1985) 387-405.
- Dodd, F.J., Donegan, H.A. and McMaster, T.B.M., A statistical approach to consistency in AHP, *Mathematical and Computer Modelling* 18 (1993) 19-22.
- Dodd, F.J., Donegan, H.A. and McMaster, T.B.M., Inverse inconsistency in analytic hierarchy process, *European Journal of Operational Research* 80 (1995) 86-93.
- Duszak, Z. and Koczkodaj, W.W., Generalization of a new definition of consistency for pairwise comparisons, *Information Processing Letters* 52 (1994) 273-276.
- Forman, E.H., Random indices for incomplete pairwise comparison matrices, *European Journal of Operational Research* 48 (1990) 153-155.
- Gass, S.I. and Rapcsák, T., Singular value decomposition in AHP, *European Journal of Operational Research* 154 (3) (2004) 573-584.
- Gass, S.I. and Standard, S.M., Characteristics of positive reciprocal matrices in the analytic hierarchy process, *Journal of Operational Research Society* 53 (2002) 1385-1389.
- Golden, B.L. and Wang, Q., An alternative measure of consistency, in: *Analytic Hierarchy Process: Applications and Studies*, B.L. Golden, E.A. Wasil and P.T. Hacker (eds.), *Springer-Verlag* (1990) 68-81.
- Jensen, R.E., Comparison of eigenvector, least squares, Chi square and logarithmic least squares methods of scaling a reciprocal matrix, *Trinity University*, Working Paper 127 (1983). (<http://www.trinity.edu/rjensen/127wp/127wp.htm>)
- Johnson, C.R., Beine, W.B. and Wang, T.J., Right-left asymmetry in an eigenvector ranking procedure, *Journal of Mathematical Psychology* 19 (1979) 61-64.
- Koczkodaj, W.W., A new definition of consistency of pairwise comparisons, *Mathematical and Computer Modelling* 8 (1993) 79-84.
- Koczkodaj, W.W., Herman, M.W. and Orlowski, M., Using consistency-driven pairwise comparisons in knowledge-based systems, in: *Proceedings of the sixth international conference on Information and knowledge management*, *ACM Press* (1997) 91-96.

- Lane, E.F. and Verdini, W.A., A consistency test for AHP decision makers, *Decision Sciences* 20 (1989) 575-590.
- Monsuur, H., An intrinsic consistency threshold for reciprocal matrices, *European Journal of Operational Research* 96 (1996) 387-391.
- Murphy, C.K., Limits on the analytic hierarchy process from its consistency index, *European Journal of Operational Research* 65 (1993) 138-139.
- Peláez, J.I. and Lamata, M.T., A new measure of consistency for positive reciprocal matrices, *Computers and Mathematics with Applications* 46 (2003) 1839-1845.
- Saaty, T.L., The analytic hierarchy process, *McGraw-Hill*, New York, 1980 and 1990.
- Saaty, T.L., Fundamentals of Decision Making, *RSW Publications*, 1994.
- Standard, S.M., Analysis of positive reciprocal matrices, Master's Thesis, Graduate School of the University of Maryland, 2000.
- Stein, W.E. and Mizzi, P.J., The harmonic consistency index for the analytic hierarchy process, *European Journal of Operational Research* 177 (2007) 488-497.
- Thorndike, E.L., A constant error in psychological ratings, *Journal of Applied Psychology* 4 (1920) 25-29.
- Thurstone, L.L., The method of paired comparisons for social values, *Journal of Abnormal and Social Psychology* 21 (1927) 384-400.
- Tummala, V.M.R. and Ling, H., A note on the computation of the mean random consistency index of the analytic hierarchy process (AHP), *Theory and Decision* 44 (1998) 221-230.
- Vargas, L.G., Reciprocal matrices with random coefficients, *Mathematical Modelling* 3 (1982) 69-81.

List of Figures 3




Figure 3.a λ_{\max} and CR values
of 3×3 random matrices




Figure 3.b λ_{\max} and CR values
of 4×4 random matrices




Figure 3.c λ_{\max} and CR values
of 5×5 random matrices




Figure 3.d λ_{\max} and CR values
of 6×6 random matrices




Figure 3.e λ_{max} and CR values of 7×7 random matrices




Figure 3.f λ_{max} and CR values of 8×8 random matrices




Figure 3.g λ_{max} and CR values of 9×9 random matrices




Figure 3.h λ_{max} and CR values of 10×10 random matrices